

A DYNAMICAL SYSTEMS APPROACH TO THE TILTED BIANCHI MODELS OF SOLVABLE TYPE

ALAN COLEY, SIGBJØRN HERVIK

ABSTRACT. We use a dynamical systems approach to analyse the tilting spatially homogeneous Bianchi models of solvable type (e.g., types VI_h and VII_h) with a perfect fluid and a linear barotropic γ -law equation of state. In particular, we study the late-time behaviour of tilted Bianchi models, with an emphasis on the existence of equilibrium points and their stability properties. We briefly discuss the tilting Bianchi type V models and the late-time asymptotic behaviour of irrotational Bianchi VII_0 models. We prove the important result that for non-inflationary Bianchi type VII_h models vacuum plane-wave solutions are the only future attracting equilibrium points in the Bianchi type VII_h invariant set. We then investigate the dynamics close to the plane-wave solutions in more detail, and discover some new features that arise in the dynamical behaviour of Bianchi cosmologies with the inclusion of tilt. We point out that in a tiny open set of parameter space in the type IV model (the loop-hole) there exists closed curves which act as attracting limit cycles. More interestingly, in the Bianchi type VII_h models there is a bifurcation in which a set of equilibrium points turn into closed orbits. There is a region in which both sets of closed curves coexist, and it appears that for the type VII_h models in this region the solution curves approach a compact surface which is topologically a torus.

1. INTRODUCTION

Spatially homogeneous (SH) models are very important in contemporary cosmology for two main reasons. First, these models are useful in analyzing the physical effects of the Universe which are affected by anisotropy in the rate of expansion. Second, SH Bianchi universes are relatively straightforward to analyse since the Einstein field equations for these cosmologies reduce to ordinary differential equations (DEs).

A SH cosmology is said to be *tilted* if the fluid velocity vector is not orthogonal to the group orbits, otherwise the model is said to be *non-tilted* [1]. Tilted universes are of considerable interest since they contain many features of physical relevance, including rotation, the conditions needed for the appearance of closed time-like curves and some properties of quantum cosmologies. In this paper we shall assume that the matter content of the universe is a perfect fluid with equation of state $p = (\gamma - 1)\mu$, where γ is constant, which includes the important cases of dust and radiation, $\gamma = 1$ and $\gamma = \frac{4}{3}$, respectively.

Much work has been done in analyzing non-tilting SH models [2, 3, 4]. However, less is known about the tilted models. Since Bianchi models with a *tilted* fluid [1]

Date: February 7, 2008.

Key words and phrases. Spatially homogeneous cosmological models, Bianchi models, dynamical systems approach, asymptotics, plane waves.

contain up to three additional degrees of freedom, it is evident that there is an increase in dynamical complexity and new features emerge with the inclusion of tilt.

For the class of tilted SH models, the Einstein field equations have been written as an autonomous DE in a number of different ways [5, 6, 7, 8, 9], but due to the complexity of the equations a detailed analysis of the dynamics is difficult. A subclass of models of Bianchi type V [10, 11, 12, 13] and models of Bianchi type II [14] have been studied using dynamical systems theory. In particular, a complete description of the dynamics of tilted SH cosmologies of Bianchi type II was given in [14], with particular emphasis on the qualitative analysis of the dynamics near the initial singularity and at late times. It was found that for $\gamma < 2$, the tilt destabilizes the Kasner solutions leading to a Mixmaster-like initial singularity, with the tilt being dynamically significant, while at late times the tilt becomes dynamically negligible unless $\gamma > \frac{10}{7}$. It was also found that the tilt does not destabilize the flat Friedmann model, with the result that the presence of tilt increases the likelihood of intermediate isotropization, supporting earlier results found in tilted Bianchi V models [13].

Recently more general tilting Bianchi type universes have been studied, with particular emphasis on exact solutions of the Einstein equations which are attractors for more general classes of solution. In [15] a local stability analysis of a wide range of tilting SH universes, including Bianchi class B models and the type VI₀ model of class A, were studied to elucidate their late-time behaviours. The asymptotic late-time behaviour of general tilted Bianchi type VI₀ universes was analysed in detail in [16], and this was generalized to Bianchi type VI₀ cosmological models containing two tilted γ -law perfect fluids in [17]. All of the future stable equilibrium points for various subclasses of tilted type VI₀ models, as well as for the general tilted type VI₀ models, were found (and all of the bifurcation values identified; e.g., $\gamma = 2/3$, $\gamma = 10/9$, $\gamma = 6/5$ and $\gamma = 4/3$).

In this paper we will generalize previous work and use a dynamical systems approach to analyse the tilting Bianchi type VI_h and VII_h models (and their subclasses) containing a γ -law perfect fluid. We shall adopt the formalism employed in [16]. These Bianchi models are of general measure in the space of all SH models and are sufficiently general to account for many interesting phenomena. In particular, the Bianchi type VII_h models are the most general models that contain the open Friedmann-Robertson-Walker (FRW) models and are consequently of special interest.

The paper is organised as follows. In the next section we shall write down the equations of motion using the orthonormal frame formalism and discuss the invariant subspaces. Various gauge choices are considered, and it is emphasised that the 'N-gauge' is useful for studying the stability of the plane wave solutions. In the next section we briefly discuss models of type V and VII₀. In particular we show that the self-similarity breaking which occurs in the non-tilted type VII₀, is also present for tilted, irrotational, type VII₀ models.

We then study the late-time behaviour of various tilted Bianchi models, with an emphasis on the existence of equilibrium points and their stability properties. In particular, we study the stability properties of the vacuum plane-wave spacetimes, and show that for $2/3 < \gamma < 2$ there will always be future stable plane-wave solutions in the set of type IV and VII_h tilted Bianchi models. Indeed, we prove

a theorem concerning the future evolution in the invariant type V and type VII_h sets, and in the case of Bianchi type VII_h models it is shown that the only future attracting equilibrium points for non-inflationary fluids ($\gamma > 2/3$) are the plane-wave solutions. We also study the form of the asymptotic tilt velocity for these attractors. For inflationary type fluids ($0 < \gamma < 2/3$) the situation is simpler; we show that the cosmic no-hair theorem is valid in this case.

We then study the dynamics in the neighbourhood of the vacuum plane-wave solutions. In the type IV models, we discover that there is a tiny region of the parameter space in which there exists a closed orbit, which acts as an attracting limit cycle. Moreover, for the type VII_h models more attracting closed orbits appear. A numerical analysis which is presented in a companion paper [18], shows that there is an open set of parameter space such that the solution curves approach a compact surface which is topologically a torus in the type VII_h models.

We also study the stability of the plane waves in Bianchi type VI_h models and the stability of the Collinson-French vacuum solution in the exceptional Bianchi type $VI_{-1/9}$ models. In the final section we summarize our results, discuss possible early time behaviour and briefly describe future work. Some more technical results are presented in the appendices.

Finally, as stressed in [3, 4], an important mathematical link exists between the various classes in the *state space* hierarchy. The physical state of a cosmological model at an instant of time is represented by a point in the state space, which is finite dimensional for SH models and infinite dimensional otherwise. This structure opens the possibility that the evolution of a model in one class may be approximated, over some time interval, by a model in a more special class. Thus it is plausible that understanding the dynamics at one level of complexity, such as for example the tilting Bianchi type VI_h and VII_h models, will shed light on the possible dynamical behaviour at a higher level, such as special classes of inhomogeneous cosmological models.

2. EQUATIONS OF MOTION

Let us consider the Bianchi models containing a tilted γ -law perfect fluid. We will investigate the models having an Abelian G_2 subgroup. For the Bianchi models, these correspond to the Lie algebras, $\mathcal{A}$, where $\mathcal{A}$ is *solvable*¹; i.e., types I- VII_h . Types VIII and IX are both semi-simple and have no Abelian G_2 subgroup. For the Bianchi cosmologies – which admit a simply transitive symmetry group acting on the spatial hypersurfaces – we can always write the line-element as

$$ds^2 = -dt^2 + \delta_{ab}\omega^a\omega^b,$$

where ω^a is a triad of one-forms obeying

$$d\omega^a = -\frac{1}{2}C^a_{bc}\omega^b\wedge\omega^c,$$

and C^a_{bc} depend only on time and are the structure constants of the Bianchi group type under consideration. The structure constants C^a_{bc} can be split into a vector part a_b , and a trace-free part n^{ab} by [2]

$$C^a_{bc} = \varepsilon_{bcd}n^{da} - \delta^a_b a_c + \delta^a_c a_b.$$

¹A Lie algebra, $\mathfrak{g}$, is called *solvable* if its derived series, defined as $\mathfrak{g}_0 = \mathfrak{g}$, $\mathfrak{g}_i = [\mathfrak{g}_{i-1}, \mathfrak{g}_{i-1}]$, terminates; i.e., there exists a k such that $\mathfrak{g}_k = 0$.

The matrix n^{ab} is symmetric, and, using the Jacobi identity, $a_b = (1/2)C_{ba}^a$ is in the kernel of n^{ab}

$$n^{ab}a_b = 0.$$

The time-like vector $\mathbf{e}_0$ is chosen orthogonal to the group orbits and is given by $\mathbf{e}_0 = \partial/\partial t$ where t is the cosmological time. It is necessary to introduce a dimensionless time variable, τ , defined by

$$(2.1) \quad \frac{dt}{d\tau} = \frac{1}{H},$$

where H is the Hubble scalar. Following [3, 4] we introduce expansion-normalised variables and the equations of motion can be given as an autonomous system of DEs (see [19]).

The equations of motion will in general depend on the choice of gauge. We will choose an orthonormal frame where $\mathbf{e}_1$ points in the direction of the vector $a_b = \frac{1}{2}C_{ba}^a$, but we will leave the remaining frame vectors $\mathbf{e}_2$ and $\mathbf{e}_3$ defined up to a rotation. Explicitly, this frame rotation is given by

$$(2.2) \quad \tilde{\mathbf{e}}_2 = \cos \phi \mathbf{e}_2 + \sin \phi \mathbf{e}_3, \quad \tilde{\mathbf{e}}_3 = -\sin \phi \mathbf{e}_2 + \cos \phi \mathbf{e}_3,$$

where ϕ is a real function. This frame rotation defines the remaining gauge freedom we have in our system of equations.

Following the notation in [19], the expansion-normalised variables are now

$$(2.3) \quad \begin{aligned} \Sigma_{ab} &= \begin{bmatrix} -2\Sigma_+ & \sqrt{3}\Sigma_{12} & \sqrt{3}\Sigma_{13} \\ \sqrt{3}\Sigma_{12} & \Sigma_+ + \sqrt{3}\Sigma_- & \sqrt{3}\Sigma_{23} \\ \sqrt{3}\Sigma_{13} & \sqrt{3}\Sigma_{23} & \Sigma_+ - \sqrt{3}\Sigma_- \end{bmatrix}, \\ N_{ab} &= \sqrt{3} \begin{bmatrix} 0 & 0 & 0 \\ 0 & \bar{N} + N_- & N_{23} \\ 0 & N_{23} & \bar{N} - N_- \end{bmatrix}. \end{aligned}$$

To facilitate the remaining gauge freedom we will utilize complex variables (which will be written in bold typeface) :

$$(2.4) \quad \begin{aligned} \mathbf{N}_\times &= N_- + iN_{23}, & \boldsymbol{\Sigma}_\times &= \Sigma_- + i\Sigma_{23} \\ \boldsymbol{\Sigma}_1 &= \Sigma_{12} + i\Sigma_{13}, & \mathbf{v} &= v_2 + iv_3. \end{aligned}$$

A gauge transformation is then given by

$$(2.5) \quad (\mathbf{N}_\times, \boldsymbol{\Sigma}_\times, \boldsymbol{\Sigma}_1, \mathbf{v}) \mapsto (e^{2i\phi}\mathbf{N}_\times, e^{2i\phi}\boldsymbol{\Sigma}_\times, e^{i\phi}\boldsymbol{\Sigma}_1, e^{i\phi}\mathbf{v}).$$

Here, the phase ϕ can be a function of time and corresponds to a rotation of frame according to eq.(2.2). From the above we see that the variables $\mathbf{N}$ and $\boldsymbol{\Sigma}_\times$ are spin-2 objects and $\boldsymbol{\Sigma}_1$ and $\mathbf{v}$ are spin-1 objects under the frame rotations. The complex conjugates, which we will denote with an asterisk, transform similarly:

$$(2.6) \quad (\mathbf{N}_\times^*, \boldsymbol{\Sigma}_\times^*, \boldsymbol{\Sigma}_1^*, \mathbf{v}^*) \mapsto (e^{-2i\phi}\mathbf{N}_\times^*, e^{-2i\phi}\boldsymbol{\Sigma}_\times^*, e^{-i\phi}\boldsymbol{\Sigma}_1^*, e^{-i\phi}\mathbf{v}^*).$$

In the following, we will explicitly leave the gauge function ϕ in the equations of motion and we will assume that ϕ is given with respect to a frame for which $\phi' = 0$ implies $R_1 = 0$ where R_a is the (expansion-normalised) local angular velocity of a Fermi-propagated axis with respect to the triad $\mathbf{e}_a$. Thus we will replace the function R_1 with the function ϕ . Explicitly, we have set $R_1 = -\phi'$. At this stage it is important to note that all the physical variables have to be independent of the gauge. Hence, the objects $\mathbf{N}_\times$ and $\mathbf{v}$ are themselves not physical variables; only

scalars constructed from the above are gauge independent (we will define what we mean by scalars below).

For the expansion-normalised variables the equations of motion are:

$$(2.7) \quad \Sigma'_+ = (q-2)\Sigma_+ + 3|\Sigma_1|^2 - 2|\mathbf{N}_\times|^2 + \frac{\gamma\Omega}{2G_+}(-2v_1^2 + |\mathbf{v}|^2)$$

$$(2.8) \quad \Sigma'_\times = (q-2+2i\phi')\Sigma_\times + \sqrt{3}\Sigma_1^2 - 2\mathbf{N}_\times(iA + \sqrt{3}\bar{N}) + \frac{\sqrt{3}\gamma\Omega}{2G_+}\mathbf{v}^2$$

$$(2.9) \quad \Sigma'_1 = (q-2-3\Sigma_+ + i\phi')\Sigma_1 - \sqrt{3}\Sigma_\times\Sigma_1^* + \frac{\sqrt{3}\gamma\Omega v_1}{G_+}\mathbf{v}$$

$$(2.10) \quad \mathbf{N}'_\times = (q+2\Sigma_+ + 2i\phi')\mathbf{N}_\times + 2\sqrt{3}\Sigma_\times\bar{N}$$

$$(2.11) \quad \bar{N}' = (q+2\Sigma_+)\bar{N} + 2\sqrt{3}\text{Re}(\Sigma_\times^*\mathbf{N}_\times)$$

$$(2.12) \quad A' = (q+2\Sigma_+)A.$$

The equations for the fluid are

$$(2.13) \quad \Omega' = \frac{\Omega}{G_+} \left\{ 2q - (3\gamma - 2) + 2\gamma Av_1 + [2q(\gamma - 1) - (2 - \gamma) - \gamma\mathcal{S}]V^2 \right\}$$

$$(2.14) \quad v'_1 = (T + 2\Sigma_+)v_1 - 2\sqrt{3}\text{Re}(\Sigma_1\mathbf{v}^*) - A|\mathbf{v}|^2 + \sqrt{3}\text{Im}(\mathbf{N}_\times^*\mathbf{v}^2)$$

$$(2.15) \quad \mathbf{v}' = \left(T - \Sigma_+ + i\phi' + Av_1 - i\sqrt{3}\bar{N}v_1 \right) \mathbf{v} - \sqrt{3}(\Sigma_\times + i\mathbf{N}_\times v_1) \mathbf{v}^*$$

$$(2.16) \quad V' = \frac{V(1-V^2)}{1-(\gamma-1)V^2} [(3\gamma-4) - 2(\gamma-1)Av_1 - \mathcal{S}]$$

where

$$(2.17) \quad \begin{aligned} q &= 2\Sigma^2 + \frac{1}{2} \frac{(3\gamma-2) + (2-\gamma)V^2}{1+(\gamma-1)V^2} \Omega \\ \Sigma^2 &= \Sigma_+^2 + |\Sigma_\times|^2 + |\Sigma_1|^2 \\ \mathcal{S} &= \Sigma_{ab}c^ac^b, \quad c^ac_a = 1, \quad v^a = Vc^a, \\ V^2 &= v_1^2 + |\mathbf{v}|^2, \\ G_+ &= 1 + (\gamma-1)V^2, \\ T &= \frac{[(3\gamma-4) - 2(\gamma-1)Av_1](1-V^2) + (2-\gamma)V^2\mathcal{S}}{1-(\gamma-1)V^2}. \end{aligned}$$

These variables are subject to the constraints

$$(2.18) \quad 1 = \Sigma^2 + A^2 + |\mathbf{N}_\times|^2 + \Omega$$

$$(2.19) \quad 0 = 2\Sigma_+A + 2\text{Im}(\Sigma_\times^*\mathbf{N}_\times) + \frac{\gamma\Omega v_1}{G_+}$$

$$(2.20) \quad 0 = \Sigma_1(i\bar{N} - \sqrt{3}A) + i\Sigma_1^*\mathbf{N}_\times + \frac{\gamma\Omega\mathbf{v}}{G_+}$$

$$(2.21) \quad 0 = A^2 + 3h(|\mathbf{N}_\times|^2 - \bar{N}^2)$$

The parameter γ will be assumed to be in the interval $\gamma \in (0, 2)$.

3. THE STATE SPACE

The state vector can be considered as $\mathbf{X} = (\Sigma_+, \Sigma_\times, \Sigma_1, \mathbf{N}_\times, \bar{N}, A, v_1, \mathbf{v})$ modulo the constraints eqs. (2.19), (2.20) and (2.21). The latter is the group constraint,

which we assume determines the parameter h and consequently is not a free variable. The Bianchi identities ensure that the constraints (2.18)-(2.20) are first integrals; thus satisfying the constraints on a initial hypersurface is sufficient to ensure that they are satisfied at all times (see also the Appendix where the expressions for their time derivatives are explicitly given).

The variable Ω is determined from the Hamiltonian constraint, eq.(2.18), and can thus be eliminated. The tuple $\mathbf{X}$ is 12-dimensional, but the constraints reduce the number of independent components to 8. In the above equations the function ϕ carries the choice of gauge and does not have an evolution equation. Specifying this function determines the gauge completely. We can use this function to eliminate one degree of freedom. This reduces the dimension of the state space to 7 (for a given h); i.e., the physical state space is 7-dimensional. Again we note that all of the physical information can be extracted from considering real scalars.

Let us list some relevant invariant subspaces necessary for this discussion.

- (1) $T(\mathcal{A})$ The full state space of tilted solvable Bianchi models.
- (2) $T(VI_h)$ Type VI_h : $|\mathbf{N}_\times|^2 - \bar{N}^2 > 0$.
- (3) $T(VII_h)$ Type VII_h : $|\mathbf{N}_\times|^2 - \bar{N}^2 < 0$.
- (4) $T(VI_0)$ Type VI_0 : $|\mathbf{N}_\times|^2 - \bar{N}^2 > 0$, $A = 0$.
- (5) $T(VII_0)$ Type VII_0 : $|\mathbf{N}_\times|^2 - \bar{N}^2 < 0$, $A = 0$.
- (6) $T(V)$ Type V : $|\mathbf{N}_\times| = \bar{N} = 0$, $A \neq 0$.
- (7) $T(IV)$ Type IV : $|\mathbf{N}_\times|^2 - \bar{N}^2 = 0$, $A \neq 0$.
- (8) $T(II)$ Type II : $|\mathbf{N}_\times|^2 - \bar{N}^2 = 0$, $A = 0$.
- (9) $B(I)$ Type I : $|\mathbf{N}_\times| = \bar{N} = A = 0$.
- (10) $T_1(\mathcal{A})$ Non-exceptional, one-component tilted fluid: $\Sigma_1 = \mathbf{v} = 0$.
- (11) $B(\mathcal{A})$ Non-tilted, non-exceptional: $\Sigma_1 = \mathbf{v} = v_1 = 0$.
- (12) $\partial T(\mathcal{A})$ “Tilted” vacuum boundary: $\Omega = 0$.
- (13) $\partial B(\mathcal{A})$ Non-tilted vacuum boundary: $\Omega = \mathbf{v} = v_1 = 0$.

Note that the Hamiltonian constraint, ensures that

$$(3.1) \quad \Sigma_+^2 + |\Sigma_\times|^2 + |\Sigma_1|^2 + |\mathbf{N}_\times|^2 + A^2 \leq 1.$$

In addition we will require that the tilt velocities are not superluminal; i.e.

$$(3.2) \quad v_1^2 + |\mathbf{v}|^2 \leq 1.$$

The variable $\bar{N}$ may or may not be bounded. However, from eq.(2.21) we see that $\bar{N}$ can only be unbounded for $T(VII_0)$. Hence, for all models except type VII_0 , the state space is compact.

3.1. Equilibrium points and gauge independent quantities.

Definition 3.1 (Scalar). We will call a quantity which is independent of the transformation, eq. (2.2), a scalar. Scalars are thus *gauge independent quantities*.

Examples of such scalars are easy to construct. Note that Σ_+ , $\bar{N}$ and A are all scalars. Furthermore, objects like $\mathbf{N}_\times^* \mathbf{N}_\times$, $\mathbf{N}_\times^* \mathbf{v}^2$, $\Sigma_1 \mathbf{v}^*$, are also scalars. In the above formalism it is also necessary to provide a gauge independent definition of equilibrium points.

Definition 3.2 (Equilibrium points). A set P is said to be (an) equilibrium point(s) if all scalars are constants on P as functions of τ .

We note that if different points in P have different scalars then the points may represent different physical configurations. The constancy as a function of τ is then the usual requirement for equilibrium points.

For the non-tilted, non-exceptional models, $\Sigma_1 = \mathbf{v} = 0$, and thus from the gauge dependent quantities $\Sigma_\times$ and $\mathbf{N}_\times$ there are only four real scalars. These are related via the identity

$$(3.3) \quad |\mathbf{N}_\times|^2 |\Sigma_\times|^2 = [\text{Re}(\mathbf{N}_\times \Sigma_\times^*)]^2 + [\text{Im}(\mathbf{N}_\times \Sigma_\times^*)]^2.$$

Hence, there are only three independent scalars constructed from $\Sigma_\times$ and $\mathbf{N}_\times$. In the non-tilted analysis, Hewitt and Wainwright [20] used these scalars as dynamical variables (instead of $\Sigma_\times$ and $\mathbf{N}_\times$ themselves) at the cost of obtaining the additional constraint (3.3). In principle, we could follow the same approach for the tilted analysis. This choice might be appropriate for certain subspaces (like $T_1(\mathcal{A})$); however, in general, this would lead to too many constraints (similar to eq.(3.3)) to be useful in practice.

3.2. Some Choices of Gauge. There are different convenient choices of gauges for the different invariant subspaces. Here we will discuss three such choices which are useful for different purposes.

3.2.1. "F-gauge": $\phi' = 0$. The simplest possible choice of ϕ is to let $\phi' = 0$. Note that this does not specify the gauge completely; we still have a constant $U(1)$ gauge freedom left. However, the F-gauge is convenient for several reasons. First, the system of equations are defined everywhere on the state space. Second, the equations of motion reduce to an autonomous system of equations which define a dynamical system (with 8 variables). However, in this gauge there is still a constant $U(1)$ -transformation remaining which can, for example, be used for simplifying the initial data. The initial data will then, modulo this constant transformation, define a unique path in the state space.

It should also be noted that in this gauge the equilibrium points of the dynamical system do not necessarily have $\mathbf{X}' = 0$. Hence, for calculating equilibrium points there might be a more convenient gauge choice.

3.2.2. "N-gauge": $\mathbf{N}_\times$ *purely imaginary*. This choice of gauge uses the gauge-function ϕ to simplify the function $\mathbf{N}_\times$. We can, for example, choose the function $\mathbf{N}_\times$ to be purely imaginary. To ensure that $\mathbf{N}_\times$ is purely imaginary at all times we have to choose $\phi' = \sqrt{3}\lambda\Sigma_-$, where λ is defined by

$$(3.4) \quad \bar{N} = \lambda \text{Im}(\mathbf{N}_\times).$$

The equation for $\bar{N}$ can then be replaced with an equation for λ which ensures a closed system of equations. The advantages of this gauge is that there is no gauge freedom left and the equilibrium points are given by $\mathbf{X}' = 0$. However, care is needed because the parameter λ is ill-defined near $\mathbf{N}_\times = 0$. Close to $\mathbf{N}_\times = 0$ the parameter λ will diverge and the dynamical system will cease to be well defined. Note, however, that in $T(VI_h)$ this cannot happen since $\lambda^2 < 1$ in $T(VI_h)$. However, in $T(VII_h)$ $\mathbf{N}_\times = 0$ may occur. If one wants to find the equilibrium points and investigate their stability then this choice of gauge is particularly useful. Equilibrium points in the neighbourhood of $\mathbf{N}_\times = 0$ have to be treated separately.

3.2.3. “ $\Sigma_1/\mathbf{v}$ -gauge”: $\Sigma_1/\mathbf{v}$ purely real. These choices of gauge are related and use the function ϕ to simplify the function Σ_1 or $\mathbf{v}$. For example, we can choose $\text{Im}(\Sigma_1) = 0$ or $\text{Im}(\mathbf{v}) = 0$. Both of these choices can be achieved by choosing ϕ to obey

$$(3.5) \quad \phi' = \sqrt{3} [\Sigma_{23} + (\bar{N} + N_-)v_1].$$

The remaining constant ϕ transformation can then be used to choose either Σ_1 or $\mathbf{v}$ purely real (however, in general both of them cannot be purely real). For this gauge, all of the variables are bounded and well-behaved. However, for $T_1(\mathcal{A})$ the gauge choice becomes degenerate so that there is still an undetermined gauge for this invariant subspace. We observe that this invariant subspace contains all of the non-tilted and the vacuum models; hence, for equilibrium points on the vacuum boundary, or non-tilted ones, this choice of gauge is not very well suited. However, in $T(\mathcal{A}) - T_1(\mathcal{A})$, this choice of gauge completely fixes the gauge and is well-behaved; this choice was used in the tilted type II analysis [14].

3.3. Equilibrium points in class B spacetimes. A number of important equilibrium points are given in the text below. Equilibrium points in the invariant Bianchi type VI_h have been studied in ref. [21]. This case contains many equilibrium points and we shall only present the plane-wave equilibrium points explicitly here (see section 5.3).

The following result is useful in the analysis of the evolution in the invariant type VII_h set, $T(\text{VII}_h)$, and the type V set, $T(V)$:

Theorem 3.3 (Type V and VII_h equilibrium points). *All equilibrium points of type V ($\mathbf{N}_\times = \bar{N} = 0$, $A \neq 0$) and VII_h ($|\mathbf{N}_\times|^2 - \bar{N}^2 < 0$, $A \neq 0$) have $\Sigma_1 = \mathbf{v} = 0$; i.e., all equilibrium points of type V and VII_h lie in the invariant subspaces $T_1(V)$ and $T_1(\text{VII}_h)$, respectively. Furthermore, equilibrium points in the interior of $T_1(\text{VII}_h)$ are either plane-wave spacetimes or FRW universes.*

Proof. The type V and VII_h cases need to be treated separately.

Type V: Assuming $\mathbf{v} \neq 0$ leads, after a fairly straightforward calculation, to a contradiction. Thus we must have $\Sigma_1 = \mathbf{v} = 0$.

Type VII_h : The proof is somewhat lengthy and an outline of the proof is given in Appendix B.1. $\square$

For $\gamma > 2/3$ the (non-vacuum) FRW universes are unstable. Therefore we immediately have:

Corollary 3.4. *In the case of Bianchi type VII_h models the only future attracting equilibrium points for non-inflationary fluids ($\gamma > 2/3$) are the plane-wave solutions.*

We also note that for inflationary fluids, $0 < \gamma < 2/3$, the FRW universe ($\Omega = 1$, $V = 0$) is stable in $T(\mathcal{A})$. In fact, for $0 < \gamma \leq 6/7$ (!) there exists a monotonically increasing function Z_1 , defined by

$$(3.6) \quad \begin{aligned} Z_1 &\equiv \alpha \Omega^{1-\gamma}, \quad \alpha = \frac{(1 - V^2)^{\frac{1}{2}(2-\gamma)}}{G_+^{1-\gamma} V^\gamma}, \\ Z_1' &= [2(1 - \gamma)q + (2 - \gamma) + \gamma\mathcal{S}] Z_1. \end{aligned}$$

To see that this is a monotonically increasing function, note first that $|\mathcal{S}| \leq 2\Sigma$ [16]. Then, using the constraint equation (2.18) and $(1 - \Sigma) \geq (1 - \Sigma^2)/2$, we can write

$$(3.7) \quad \begin{aligned} & 2(1 - \gamma)q + (2 - \gamma) + \gamma\mathcal{S} \\ & \geq (6 - 7\gamma)\Sigma^2 + 2(1 - \gamma)(|\mathbf{N}_\times|^2 + A^2) + \frac{\gamma(1 - \gamma)(V^2 + 3)}{G_+}\Omega, \end{aligned}$$

which is strictly positive for $\gamma < 6/7$. Thus for $0 < \gamma \leq 6/7$ Z_1 is monotonically increasing as claimed.

Theorem 3.5. *For $0 < \gamma \leq 6/7$, all tilted Bianchi models (with $\Omega > 0$, $V < 1$) of solvable type are asymptotically non-tilted at late times.*

Proof. Use of the monotonic function Z_1 . □

In fact, the result that these models are asymptotically non-tilted at late times is true for all $\gamma < 1$; this follows from an analysis, but is not covered by this theorem. This theorem is, in fact, true for all ever-expanding Bianchi models, including ever-expanding class A models².

An immediate result of the above is the following corollary:

Corollary 3.6 (Cosmic no-hair). *For $\Omega > 0$, $V < 1$, and $0 < \gamma < 2/3$ we have that*

$$\lim_{\tau \rightarrow \infty} \Omega = 1, \quad \lim_{\tau \rightarrow \infty} V = 0.$$

Proof. Using Theorem 3.5 we can write

$$\Omega' = \Omega [2q - (3\gamma - 2) + \delta]$$

where $\delta \rightarrow 0$ as $\tau \rightarrow \infty$. Furthermore, for $0 < \gamma < 2/3$,

$$2q - (3\gamma - 2) \geq (2 - 3\gamma)(1 - \Omega) \geq 0,$$

where equality can only occur for $\Omega = 1$. Assuming $\Omega < 1$, then for every $0 < \gamma < 2/3$ there will exist a τ_1 such that $2q - (3\gamma - 2) + \delta > 0$ for all $\tau > \tau_1$. The corollary now follows. □

This result is usually referred to as the cosmic no-hair theorem; the original version given by Wald [22] was for a cosmological constant. The above corollary generalises the non-tilted result given in [3]. We shall be primarily interested in the future asymptotic behaviour of models with $2/3 < \gamma < 2$ henceforward.

²For class A models ($A = 0$) the upper bound $6/7$ can be slightly improved to $14/15$ by modifying the monotonic function Z_1 to

$$\tilde{Z}_1 \equiv \tilde{\alpha}\Omega^{1-\beta}, \quad \tilde{\alpha} = \frac{(1 - V^2)^{\frac{1}{2}(2-\gamma)}}{G_+^{1-\beta}V^\beta}, \quad \beta = \frac{6}{7}\gamma.$$

4. SPECIAL SUBSETS

4.1. Bianchi Type V. The simplest model of class B is the type V model for which $\mathbf{N}_\times = \bar{N} = 0$, $A \neq 0$. Hewitt and Wainwright [13] studied a subclass of this model, namely the evolution in the $T_1(V)$ invariant subspace.

For the type V models the constraint (2.20) simplifies to

$$(4.1) \quad \sqrt{3}A\Sigma_1 = \frac{\gamma\Omega}{G_+}\mathbf{v},$$

which implies that Σ_1 and $\mathbf{v}$ are parallel. Thus for type V the Σ_1 and $\mathbf{v}$ gauges are equivalent.

Theorem 3.3 implies that all of the equilibrium points of type V are given in Hewitt and Wainwright's analysis [13]. Hence, the equilibrium points can be extracted directly from that paper. The stability analysis, on the other hand, must be performed in the full state space which will change the stability properties of the equilibrium points.

The equilibrium points important for the late-time behaviour in the invariant space $T_1(V)$ are as follows:

- (1) $\mathcal{M}$, Milne: Future stable in $T_1(V)$ for $2/3 < \gamma < 4/3$.
- (2) $\widetilde{\mathcal{M}}$, "Tilted Milne": Future stable in $T_1(V)$ for $4/3 < \gamma < 2$.
- (3) $\mathcal{M}_\pm$, "Extremely tilted Milne": $\mathcal{M}_-$ is future stable in $T_1(V)$ for $6/5 < \gamma < 2$. $\mathcal{M}_+$ unstable for $0 < \gamma < 2$.

By considering these equilibrium points in the full type $T(V)$ state space we obtain the following stability properties:

- (1) $\mathcal{M}$: Future stable in $T(V)$ for $2/3 < \gamma < 4/3$.
- (2) $\widetilde{\mathcal{M}}$: Unstable in $T(V)$.
- (3) $\mathcal{M}_\pm$: $\mathcal{M}_-$ is future stable in $T(V)$ for $6/5 < \gamma < 2$. $\mathcal{M}_+$ unstable for $0 < \gamma < 2$.

As we shall see later, these results can be directly extracted from the plane wave analysis (even though we assume $\mathbf{N}_\times \neq 0$ in this case). The reason for this is that the Milne universe can be considered the isotropic limit of the vacuum plane waves ($\Sigma_+, \bar{N} \rightarrow 0$).

4.2. Irrotational Bianchi type VII₀. Let us take a first look at the tilted Bianchi type VII₀ model. For the non-tilted model it is known that the variable $\bar{N}$ diverges as $\tau \rightarrow \infty$ [23, 24] which causes the non-tilted type VII₀ models to experience a *self-similarity breaking* at late times [23, 24]. We will show that this self-similarity breaking persists within the subspace $T_1(VII_0)$. In the non-tilted case, the variable $\mathbf{N}_\times$ will oscillate around the value zero. To avoid the problem of the $\mathbf{N}$ -gauge at $\mathbf{N}_\times = 0$, we will choose the F-gauge for which $\phi' = 0$.

We will consider the invariant subspace $T_1(VII_0)$, where $\mathbf{v} = \Sigma_1 = 0$, and we will assume, with no loss of generality, that $\bar{N}, v_1 > 0$. In this invariant subspace

there are two monotonic functions,

$$\begin{aligned} Z_2 &\equiv \frac{|\mathbf{N}_\times|^2 + |\boldsymbol{\Sigma}_\times|^2}{\bar{N}^2 + |\boldsymbol{\Sigma}_\times|^2}, & \frac{Z'_2}{Z_2} &= -\frac{4|\boldsymbol{\Sigma}_\times|^2 (\bar{N}^2 - |\mathbf{N}_\times|^2) (\Sigma_+ + 1)}{(|\mathbf{N}_\times|^2 + |\boldsymbol{\Sigma}_\times|^2) (\bar{N}^2 + |\boldsymbol{\Sigma}_\times|^2)}, \\ Z_3 &\equiv \frac{(\bar{N}^2 - |\mathbf{N}_\times|^2)^n \beta \Omega}{(1 + n\Sigma_+)^{2(1+n)}}, & \frac{Z'_3}{Z_3} &= \frac{4 \left[(\Sigma_+ + n)^2 + (1 - n^2) \left(|\boldsymbol{\Sigma}_\times|^2 + \frac{(1+2n)\Omega V^2}{3G_+} \right) \right]}{1 + n\Sigma_+}, \end{aligned} \quad (4.2)$$

where

$$(4.3) \quad \beta \equiv \frac{(1 - V^2)^{\frac{1}{2}(2-\gamma)}}{G_+}, \quad n \equiv \frac{1}{4}(3\gamma - 2).$$

Theorem 4.1. *For a non-inflationary perfect fluid ($2/3 < \gamma < 2$) and any initial value in $T_1(VII_0)$ with $V < 1$ and $\Omega > 0$, we have that*

$$\lim_{\tau \rightarrow +\infty} |\bar{N}| = \infty.$$

Proof. Use of the monotonic function Z_3 (the proof is completely analogous to the non-tilted case [23]). $\square$

We note that this implies that these models are not asymptotically self-similar at late-times. Let us thus assume that $1/\bar{N} \rightarrow 0$ and investigate what consequences this may have for the evolution of the tilted type VII₀ model. For $\bar{N} \rightarrow \infty$, the equations for $\mathbf{N}_\times$ and $\boldsymbol{\Sigma}_\times$ have the asymptotic form

$$(4.4) \quad \begin{aligned} \mathbf{N}'_\times &= 2\sqrt{3}\bar{N}\boldsymbol{\Sigma}_\times, \\ \boldsymbol{\Sigma}'_\times &= -2\sqrt{3}\bar{N}\mathbf{N}_\times. \end{aligned}$$

These equations can be solved in quadrature. By defining $\varphi \equiv \int_{\tau_0}^{\tau} 2\sqrt{3}\bar{N}d\tau$, we have the asymptotic form

$$(4.5) \quad \mathbf{N}_\times = \frac{1}{2}(\alpha_0 e^{-i\varphi} + \beta_0 e^{i\varphi}), \quad \boldsymbol{\Sigma}_\times = \frac{1}{2i}(\alpha_0 e^{-i\varphi} - \beta_0 e^{i\varphi}).$$

We also note that if we require $\mathbf{N}_\times \neq 0$ for all τ , then $|\alpha_0| \neq |\beta_0|$.

We have a similar type of oscillation in this case. This can be verified by calculating the scalar $|\boldsymbol{\Sigma}_\times|^2$:

$$(4.6) \quad |\boldsymbol{\Sigma}_\times|^2 = \frac{1}{4} (|\alpha_0|^2 + |\beta_0|^2 - 2|\alpha_0||\beta_0| \cos 2\varphi).$$

We also note that

$$(4.7) \quad \text{Im}(\boldsymbol{\Sigma}_\times \mathbf{N}_\times^*) = \frac{1}{4} (|\alpha_0|^2 - |\beta_0|^2),$$

so that according to the constraint (2.19), the type VII₀ model ($A = 0$) has non-zero v_1 if $|\alpha_0| \neq |\beta_0|$ in the limit $\bar{N} \rightarrow \infty$.

Based on these initial investigations, we introduce the four scalars

$$(4.8) \quad \begin{aligned} \sigma_1 &= |\mathbf{N}_\times|^2 + |\boldsymbol{\Sigma}_\times|^2, & \sigma_2 &= 2\text{Im}(\boldsymbol{\Sigma}_\times \mathbf{N}_\times^*) \\ \sigma_3 &= |\mathbf{N}_\times|^2 - |\boldsymbol{\Sigma}_\times|^2, & \sigma_4 &= 2\text{Re}(\boldsymbol{\Sigma}_\times \mathbf{N}_\times^*). \end{aligned}$$

At late times, the scalars σ_3 and σ_4 will oscillate with increasing frequency. Following [23], we incorporate this oscillation into a phase ψ by defining

$$(4.9) \quad \sigma_3 = \rho \sin \psi, \quad \sigma_4 = \rho \cos \psi.$$

Furthermore, by defining $M \equiv 1/\bar{N}$, the equations of motion can be written

$$(4.10) \quad \Sigma'_+ = (Q - 2)\Sigma_+ - \sigma_1 - \frac{\gamma\Omega V^2}{G_+} - (1 + \Sigma_+)\rho \sin \psi,$$

$$(4.11) \quad \sigma'_1 = 2(Q + \Sigma_+ - 1)\sigma_1 + (\Sigma_+ + 1 - \sigma_1)\rho \sin \psi,$$

$$(4.12) \quad V' = \frac{V(1 - V^2)}{1 - (\gamma - 1)V^2} [(3\gamma - 4) + 2\Sigma_+],$$

$$(4.13) \quad \rho' = 2[(Q + \Sigma_+ - 1)\rho + (\Sigma_+ + 1)\sigma_1 \sin \psi - \rho^2 \sin \psi],$$

$$(4.14) \quad M' = -M \left(Q + 2\Sigma_+ - \rho \sin \psi + \sqrt{3}M\rho \cos \psi \right),$$

$$(4.15) \quad \psi' = \frac{1}{M} \left[4\sqrt{3} + 2M(\Sigma_+ + 1)\frac{\sigma_1}{\rho} \cos \psi \right],$$

where

$$(4.16) \quad Q = 2\Sigma_+^2 + \sigma_1 + \frac{1}{2} \frac{(3\gamma - 2) + (2 - \gamma)V^2}{G_+} \Omega.$$

The variables Ω and σ_2 are determined from the constraints,

$$(4.17) \quad \Omega = 1 - \sigma_1 - \Sigma_+^2, \quad \sigma_2 = -\frac{\gamma\Omega V}{G_+}.$$

Moreover, the identity (3.3) implies the constraint

$$(4.18) \quad \sigma_1^2 = \sigma_2^2 + \rho^2.$$

The analysis in [23] showed that for the non-tilted model, effectively the oscillatory terms in the system of equations are dynamically negligible at late times. A similar analysis shows that the same is true in this case. This means that at late times we obtain a reduced system of equations by simply dropping the terms containing $\sin \psi$ and $\cos \psi$. This enables us to state:

Theorem 4.2 (Global attractor for $2/3 < \gamma < 4/3$). *For $2/3 < \gamma < 4/3$, a non-LRS universe in $T_1(VII_0)$ with $V < 1$ and $\Omega > 0$ has*

$$(4.19) \quad \lim_{\tau \rightarrow \infty} (\Sigma_+, \sigma_1, \rho, V) = (0, 0, 0, 0), \quad \lim_{\tau \rightarrow \infty} \Omega = 1.$$

Proof. The proof is completely analogous to that in [23] (by replacing Ω with $\beta\Omega$, where β is defined in eq.(4.3)), or using $\hat{Z}_4$ defined in Appendix B.2. $\square$

When the fluid is stiffer than radiation ($4/3 < \gamma < 2$), the universe will not isotropise in terms of the shear. By studying the reduced system of equations – which is obtained by simply dropping the terms containing sine and cosine in eqs.(4.10)–(4.15) – we can show that a solution asymptotes to a point on the following line-bifurcation:

$$(4.20) \quad \begin{aligned} \Sigma_+ &\rightarrow -\frac{1}{2}(3\gamma - 4), \quad Q \rightarrow \frac{1}{2}(3\gamma - 2), \\ \sigma_1 &\rightarrow \frac{3(2 - \gamma)}{4(1 - V^2)} [(3\gamma - 4) - V^2(5\gamma - 4)], \quad \sigma_2 \rightarrow -\frac{3V\gamma(2 - \gamma)}{2(1 - V^2)}, \\ \rho &\rightarrow \frac{3(2 - \gamma)}{4(1 - V^2)^{\frac{1}{2}}} \sqrt{(3\gamma - 4)^2 - V^2(5\gamma - 4)^2}, \quad \Omega \rightarrow \frac{3(2 - \gamma)}{2(1 - V^2)} [1 + (\gamma - 1)V^2], \\ 0 &\leq V \leq \frac{3\gamma - 4}{5\gamma - 4}. \end{aligned}$$

Theorem 4.3 (Global attractors for $4/3 < \gamma < 2$). *For $4/3 < \gamma < 2$, a non-LRS universe in $T_1(VII_0)$ with $V < 1$ and $\Omega > 0$ has*

$$(4.21) \quad \lim_{\tau \rightarrow \infty} \Sigma_+ = -\frac{1}{2}(3\gamma - 4),$$

and consequently approaches the line given by eq.(4.20).

Proof. A proof is provided in Appendix B.2. $\square$

To summarise, these results imply that for perfect fluids with $2/3 < \gamma < 4/3$ the $T_1(VII_0)$ models isotropise in terms of the shear, while for fluids with $4/3 < \gamma < 2$, the $T_1(VII_0)$ models *do not* isotropise in terms of the shear. This result is similar to the non-tilted case [23]. Regarding the expansion-normalised Weyl tensor we thus expect a similar behaviour as in the non-tilted case. In the terminology of [29] the $T_1(VII_0)$ models are, for fluids stiffer than dust ($\gamma = 1$), *extremely Weyl dominant* at late times. Note that the radiation case, $\gamma = 4/3$, has not been treated here; this case requires a separate analysis [24].

In the full tilted Bianchi VII_0 case we expect $\bar{N} \rightarrow \infty$ and a similar behaviour to occur. Tilted type VII_0 models is presently being studied by other authors [25].

5. STABILITY OF THE PLANE WAVES

In this section we will address the question: Are the vacuum plane waves stable in the set of tilted Bianchi models? To answer this question we must first define what we mean by the “vacuum plane waves”. These plane-wave solutions correspond to an invariant set, $\mathcal{P}(\mathcal{A})$, of the dynamical system. Since this set is vacuum, $\Omega = 0$, the tilt velocity decouples from the equations of motion and becomes unphysical in the invariant set itself. However, a perturbation away from the vacuum solutions generically has $\Omega \neq 0$, and in this case it is meaningful to talk about tilt. Hence, we may have a situation where the asymptotic solutions are identical but where the asymptotic tilt velocities are different.

To see how the system of equations decouples near vacuum equilibrium points, we linearise the full system of equations with respect to an equilibrium point with $\Omega = 0$. Let $\mathbf{Y}$ be a column vector corresponding to the variables given by the set of eqs. (2.7)-(2.12), and $\mathbf{v}$ the vector (v_1, v_2, v_3) . Furthermore, we assume that the constraint equations have been used to reduce the vector $\mathbf{Y}$. Then, at the linearised level, we can write

$$(5.1) \quad \begin{bmatrix} \delta \mathbf{Y} \\ \delta \Omega \\ \delta \mathbf{v} \end{bmatrix}' = \begin{bmatrix} \mathbf{A} & \lambda_{\mathbf{Y}} & 0 \\ 0 & \lambda_{\Omega} & 0 \\ \mathbf{C} & \lambda_{\mathbf{v}} & \mathbf{B} \end{bmatrix} \begin{bmatrix} \delta \mathbf{Y} \\ \delta \Omega \\ \delta \mathbf{v} \end{bmatrix}.$$

We note that the eigenvalue equation reduces to

$$(5.2) \quad \det(\mathbf{A} - \Lambda \mathbf{1}_1) \det(\mathbf{B} - \Lambda \mathbf{1}_2) (\lambda_{\Omega} - \Lambda) = 0.$$

Hence, near the vacuum equilibrium points, the system of equations decouples and we can study the two decoupled systems separately. In addition, we need to check the value of λ_{Ω} to make sure that the vacuum approximation remains valid.

Since we are dealing with an invariant set for which $\Omega = 0$, it is convenient to choose the $\mathbf{N}_{\times}$ -gauge, where the case $\lambda = \infty$ correspond to $N_{23} = 0$. We recall that in general $\bar{N} = \lambda N_{23}$.

By vacuum plane waves we will mean all solutions where:

Vacuum plane waves: $\mathcal{P}(\mathcal{A})$, $(0 < \gamma < 2)$

$$(5.3) \quad \begin{aligned} q &= -2\Sigma_+, & N_{23}^2 &= -\Sigma_+(1 + \Sigma_+), & A^2 &= (1 + \Sigma_+)^2, \\ \Sigma_-^2 &= -\Sigma_+(1 + \Sigma_+), & \Sigma_{12} &= \Sigma_{13} = \Sigma_{23} = \Omega = 0. \end{aligned}$$

The parameter h is determined by λ via

$$3h\Sigma_+(1 - \lambda^2) = (1 + \Sigma_+),$$

and Σ_+ is bounded by

$$(5.4) \quad -1 < \Sigma_+ \begin{cases} < 0, & \lambda^2 > 1 \text{ (VII}_h\text{)}, \\ < 0, & \lambda^2 = 1 \text{ (IV)}, \\ \leq \frac{1}{3h-1}, & \lambda^2 < 1 \text{ (VI}_h\text{)}. \end{cases}$$

Without loss of generality we can also assume that $A, \Sigma_-, N > 0$.

To investigate the various versions of the vacuum plane waves (i.e., with different tilt velocities) we can assume that the geometric variables have the above values and use these in the equations for v_1, v_2, v_3 . The paradigm is that as the universe approaches a plane wave, the variables Σ_+, Σ_- , etc. 'freeze in' and effectively become constants. There are several results that support this paradigm. A stability analysis shows that this is true locally. Furthermore, comprehensive numerical analysis in the full state space suggests that this is true globally; indeed, evidence for other behaviour has not been seen [18]. Once these variables are frozen, we obtain a reduced three-dimensional dynamical system that can be studied. These equations can therefore be reinterpreted as describing the tilt velocity of test matter in a plane-wave background. For convenience, and to distinguish them from the full set of equations, we define $(x, y, z) \equiv \left(v_1, \frac{v_2+v_3}{\sqrt{2}}, \frac{v_2-v_3}{\sqrt{2}}\right)$. The reduced equations for (x, y, z) are then:

$$(5.5) \quad \begin{aligned} x' &= (T + 2\Sigma_+)x - A(y^2 + z^2) - 2\sqrt{3}N_{23}yz, \\ y' &= (T - \Sigma_+ + Ax)y - \alpha(1 - \lambda)z, \\ z' &= (T - \Sigma_+ + Ax)z - \alpha(1 + \lambda)y, \end{aligned}$$

where $\alpha = \sqrt{3}(\Sigma_- - xN_{23}) = (1 - x)\sqrt{3}\Sigma_-$, since $N_{23} = \Sigma_-$ for the plane-wave solutions. The function T is given by eq.(2.17) with

$$V^2 = x^2 + y^2 + z^2, \quad V^2\mathcal{S} = \Sigma_+(-2x^2 + y^2 + z^2) + 2\sqrt{3}\Sigma_-yz.$$

The requirement $0 \leq V \leq 1$, implies

$$0 \leq x^2 + y^2 + z^2 \leq 1.$$

Hence, the state space is the solid unit ball, D^3 .

To get some sense of the resulting dynamics, we note that

$$(5.6) \quad R' = (T - \Sigma_+ + Ax)R, \quad R^2 \equiv (\lambda + 1)y^2 + (\lambda - 1)z^2.$$

Hence, $R, x = \text{constant}$ describes ellipses for type VII_h ($\lambda^2 > 1$), straight lines for type IV ($\lambda^2 = 1$), and hyperbolae for type VI_h ($\lambda^2 < 1$). (Note that since these equations are not $U(1)$ invariant, there is no question that the resulting dynamics is a gauge effect.)

Let us study the reduced dynamical system (5.5). The key observation is that from this reduced system, and from equation eq.(2.13) for Ω , we can say something about the stability of the plane waves. The strategy is as follows: We first find the

asymptotic behaviour of the system (5.5). If, in addition, the variable Ω is stable with respect to this asymptotic behaviour, then the vacuum plane waves will be stable in the full set of equations. Here, we will present heuristic arguments for some of the features of these model. In a companion paper [18] these features are more thoroughly investigated.

The primary aim is the investigation of the type VII_h plane waves; however, it is very illustrative (and also convenient) to consider the type IV plane waves first. We will also consider the type VI_h plane-waves.

5.1. Type IV plane waves. This case is defined by $\lambda = 1$, which makes R degenerate. The invariant surface $R = y = 0$, splits the solid ball, D^3 , into two equal halves which are related via the symmetry $(x, y, z) \rightarrow (x, -y, -z)$. It therefore suffices to consider the half in which $y \geq 0$. The equilibrium points are:

- (1) $\mathcal{L}(IV)$: Non-tilted, $x = y = z = 0$.
- (2) $\tilde{\mathcal{L}}(IV)$: Intermediately tilted, $x = \frac{3\gamma-4+2\Sigma_+}{2(\gamma-1)(1+\Sigma_+)}$, $y = z = 0$, $\frac{6}{5+2\Sigma_+} < \gamma < 2$.
- (3) $\tilde{\mathcal{L}}_+(IV)$: Extremely tilted, $x = 1$, $y = z = 0$.
- (4) $\tilde{\mathcal{L}}_-(IV)$: Extremely tilted, $x = -1$, $y = z = 0$.
- (5) $\tilde{\mathcal{F}}_{\pm}(IV)$: Intermediately tilted, $x = -\frac{3\gamma-4-\Sigma_+}{(1+\Sigma_+)(3-2\gamma)}$,
 $z = \pm \frac{\sqrt{(1-2\Sigma_+)(4-3\gamma)(3\gamma-4-\Sigma_+)}}{(1+\Sigma_+)(3-2\gamma)}$, $y = 0$. $\frac{4+\Sigma_+}{3} < \gamma < \min\left(\frac{4}{3}, \frac{3}{2-\Sigma_+}\right)$.
- (6) $\tilde{\mathcal{E}}_{\pm}(IV)$: Extremely tilted, $x = \frac{1+\Sigma_+}{3\Sigma_+}$, $z = \pm \frac{\sqrt{(1-2\Sigma_+)(-1-4\Sigma_+)}}{3|\Sigma_+|}$, $y = 0$,
 $-1 < \Sigma_+ < -\frac{1}{4}$.

For various different values of Σ_+ and γ these equilibrium points serve as attractors, saddle points or repellers. As we are mostly interested in the future asymptotic behaviour, we will only be concerned with the future attractors. In addition to the eigenvalues arising from the above reduced dynamical system we have to make sure that the eigenvalue from the Ω -equation is negative as well. This will ensure that $\Omega \rightarrow 0$ at late times so that our approximation is valid. Furthermore, one of the eigenvalues for $\tilde{\mathcal{F}}_{\pm}(IV)$ and $\tilde{\mathcal{E}}_{\pm}(IV)$ is zero (namely, the one in the y -direction) so we have to go to next order to determine their stability. This is just the reflection of the fact that $\tilde{\mathcal{F}}_{\pm}(IV)$ and $\tilde{\mathcal{E}}_{\pm}(IV)$ can only be an attractor in one of the half-spheres, and not in the other.

The equilibrium points are attractors for the region $y \geq 0$ for the following range of parameters:

- (1) $\mathcal{L}(IV)$: $\frac{2-4\Sigma_+}{3} < \gamma < \frac{4+\Sigma_+}{3}$.
- (2) $\tilde{\mathcal{L}}(IV)$: Always unstable for $0 < \gamma < 2$.
- (3) $\tilde{\mathcal{L}}_+(IV)$: Always unstable for $0 < \gamma < 2$.
- (4) $\tilde{\mathcal{L}}_-(IV)$: $\frac{6}{5+2\Sigma_+} < \gamma < 2$, $-\frac{1}{4} < \Sigma_+ < 0$.
- (5) $\tilde{\mathcal{F}}_{\pm}(IV)$: $\tilde{\mathcal{F}}_-(IV)$ stable³ for $\max\left(\frac{6}{5}, \frac{4+\Sigma_+}{3}\right) < \gamma < \min\left(\gamma_0, \frac{3}{2-\Sigma_+}\right)$
- (6) $\tilde{\mathcal{E}}_{\pm}(IV)$: $\tilde{\mathcal{E}}_-(IV)$ stable for $-\frac{1}{2} < \Sigma_+ < -\frac{1}{4}$, $\frac{3}{2-\Sigma_+} < \gamma < 2$.

Here, we have defined γ_0 for any given value of Σ_+ as $F(\gamma_0, \Sigma_+) = 0$, where $F(\gamma, \Sigma_+)$ is defined in eq.(C.3) in the Appendix. Note that there is a region where there are two co-existing future attractors; namely, the region where $\frac{6}{5+2\Sigma_+} < \gamma < \gamma_0$. Also, there appears to be a tiny region $\gamma_0 < \gamma < \frac{6}{5+2\Sigma_+}$ (from now on called

³ $\tilde{\mathcal{F}}_+(IV)$ is stable in the same region of γ for the half $y \leq 0$.




FIGURE 1. The regions where the different equilibrium points are attractors for the type IV model with $y \geq 0$. In the left figure, all of the boundaries along the left edge mark the instability of the energy density, Ω . The right figure is a magnified region of the left one showing the loophole.

the 'loophole') which does not have any stable equilibrium points (see Figure 1). This perhaps suggests that there is some interesting dynamical behaviour in this region. Numerical experimentation seems to indicate that there is a *stable curve*, $\mathcal{C}(IV)$, which acts as an attractor in this 'loophole'. This curve lies in the invariant subspace $y = 0$ and has a 'mussel-like' shape. In Figure 2 we have plotted two solution curves for two different values of the parameters γ and Σ_+ . From the figure one can clearly see the curve, $\mathcal{C}(IV)$, which acts as a limit cycle. Clearly it is not sufficient to only consider equilibrium points as attractors when investigating the dynamics close to a plane-wave spacetime in these SH models. These limit cycles are dealt with more elaborately in the work [18].

From the above we can conclude: For $2/3 < \gamma < 2$ there will always be future stable plane-wave spacetimes for the type IV model. Regarding the asymptotic tilt velocity, we have the following possibilities depending on the value of γ .

- $2/3 < \gamma < 6/5$: The tilt is asymptotically zero [$\mathcal{L}(IV)$]; i.e. non-tilted.
- $6/5 < \gamma < 4/3$: The tilt can be either zero [$\mathcal{L}(IV)$], intermediate [$\tilde{\mathcal{F}}(IV)$ or $\mathcal{C}(IV)$], or extreme [$\tilde{\mathcal{L}}_-(IV)$ or $\tilde{\mathcal{E}}(IV)$].
- $4/3 < \gamma < 2$: The tilt is asymptotically extreme [$\tilde{\mathcal{L}}_-(IV)$ or $\tilde{\mathcal{E}}(IV)$].

5.2. Type VII_h plane waves. In this case $\lambda^2 > 1$, and the level curves of R describe elliptic cylinders. We introduce elliptic polar coordinates by

$$(\sqrt{\lambda+1}y, \sqrt{\lambda-1}z) = (R \cos \varphi, R \sin \varphi),$$

where the evolution equation for φ is given by

$$(5.7) \quad \varphi' = (1-x)\sqrt{3(\lambda^2-1)}\Sigma_-.$$




FIGURE 2. The Mussel attractor of type IV: Two solution curves where the parameters are given inside (left), and just outside (right) the loophole. The curves are projected onto the zx -plane. Note the closed curve, $\mathcal{C}(IV)$, which acts as an attractor. On the right figure, the equilibrium point $\tilde{\mathcal{F}}_-(IV)$ acts as a future attractor. For both figures $x < 0$ and $z < 0$ at late times.

Since Σ_- and λ are constants for the plane waves, the tilt velocity will have a non-zero elliptic angular velocity with respect to the x -axis (in agreement with theorem 3.3). All of the equilibrium points must therefore lie on the x -axis:

- (1) $\mathcal{L}(h)$: Non-tilted, $x = y = z = 0$.
- (2) $\tilde{\mathcal{L}}(h)$: Intermediately tilted, $x = \frac{3\gamma-4+2\Sigma_+}{2(\gamma-1)(1+\Sigma_+)}$, $y = z = 0$, $\frac{6}{5+2\Sigma_+} < \gamma < 2$.
- (3) $\tilde{\mathcal{L}}_+(h)$: Extremely tilted, $x = 1$, $y = z = 0$.
- (4) $\tilde{\mathcal{L}}_-(h)$: Extremely tilted, $x = -1$, $y = z = 0$.

One might wonder what happened to the equilibrium points $\tilde{\mathcal{F}}_{\pm}(IV)$ and $\tilde{\mathcal{E}}_{\pm}(IV)$ as we went over to the type VII_h models. In fact, these equilibrium points have turned into *closed periodic orbits*. Hence, also in this case, it is not sufficient to consider only the equilibrium points; there are also exact closed periodic orbits that may act as future attractors. These closed periodic orbits are difficult to deal with in full generality; however, analytical, as well as numerical results concerning the existence of these orbits of the type VII_h model are presented in [18].

The stability of the type VII_h plane waves appears to be similar to that of the type IV model, except that the points $\tilde{\mathcal{F}}_{\pm}(IV)$ and $\tilde{\mathcal{E}}_{\pm}(IV)$ have become closed orbits. These closed orbits appear to be stable in approximately the same region as their IV counterparts; however, there is some uncertainty regarding the boundary of the loophole.

Regarding the closed curve $\mathcal{C}(IV)$ in the 'loophole', this seems to have turned into something more complicated for the type VII_h models. Naively, it might be anticipated that the attracting set $\mathcal{C}(IV)$ has turned into something which is topologically a torus, $\mathcal{T}$. Numerical work confirm this claim [18].

In any case we can conclude: For $2/3 < \gamma < 2$ there will always be future stable plane-wave spacetimes for the type VII_h model. Regarding the asymptotic tilt velocity, we have the following possibilities depending on the value of γ :

- $2/3 < \gamma < 6/5$: The tilt is asymptotically zero $[\mathcal{L}(h)]$; i.e. non-tilted.
- $6/5 < \gamma < 4/3$: The tilt can be either zero $[\mathcal{L}(h)]$, intermediate and oscillatory, extreme $[\tilde{\mathcal{L}}_-(h)]$, or extreme and oscillatory.
- $4/3 < \gamma < 2$: The tilt is asymptotically extreme, and can either be non-oscillatory $[\tilde{\mathcal{L}}_-(h)]$ or oscillatory.

5.3. Type VI_h plane waves. In this case we have $\lambda^2 < 1$, and the level curves of R are hyperbolae. The invariant subspace $R = 0$ defines two planes intersecting at $y = z = 0$ and divides the ball D^3 into 4 pieces. Due to the symmetry $(x, y, z) \rightarrow (x, -y, -z)$, the 4 pieces are 2 equivalent pairs. Regarding the equilibrium points, it therefore suffices to consider only equilibrium points in the region $y \geq 0$. All of the equilibrium points reside in the invariant subspace $R = 0$ (we define $\tilde{h} = 1/\sqrt{|h|}$):

- (1) $\mathcal{L}(h)$: Non-tilted, $x = y = z = 0$.
- (2) $\tilde{\mathcal{L}}(h)$: Intermediately tilted, $x = \frac{3\gamma-4+2\Sigma_+}{2(\gamma-1)(1+\Sigma_+)}$, $y = z = 0$, $\frac{6}{5+2\Sigma_+} < \gamma < 2$.
- (3) $\tilde{\mathcal{L}}_+(h)$: Extremely tilted, $x = 1$, $y = z = 0$.
- (4) $\tilde{\mathcal{L}}_-(h)$: Extremely tilted, $x = -1$, $y = z = 0$.
- (5) $\tilde{\mathcal{F}}_\pm(h)$: Intermediately tilted, $x = -\frac{3\gamma-4-\Sigma_+\mp(1+\Sigma_+)\tilde{h}}{(1+\Sigma_+)(3-2\gamma\pm\tilde{h})}$, $y = \chi\sqrt{1-\lambda}/\sqrt{2}$,
 $z = \pm\chi\sqrt{1+\lambda}/\sqrt{2}$ where

$$\chi = \frac{\sqrt{(1-2\Sigma_+)[4-3\gamma\pm\tilde{h}(2-\gamma)][3\gamma-4-\Sigma_+\mp(1+\Sigma_+)\tilde{h}]}}{(1+\Sigma_+)(3-2\gamma\pm\tilde{h})\sqrt{1\pm\tilde{h}}},$$

$$\frac{4+\Sigma_+\pm(1+\Sigma_+)\tilde{h}}{3} < \gamma < \min\left(\frac{4\pm 2\tilde{h}}{3\pm\tilde{h}}, \frac{3}{2-\Sigma_+\mp(1+\Sigma_+)\tilde{h}}\right).$$

- (6) $\tilde{\mathcal{E}}_\pm(h)$: Extremely tilted, $x = \frac{(1+\Sigma_+)(1\pm\tilde{h})}{3\Sigma_+\pm(1+\Sigma_+)\tilde{h}}$, $y = \chi\sqrt{1-\lambda}/\sqrt{2}$,
 $z = \pm\chi\sqrt{1+\lambda}/\sqrt{2}$ where

$$\chi = \frac{\sqrt{(1-2\Sigma_+)(-1-4\Sigma_+\mp 2(1+\Sigma_+)\tilde{h})}}{|3\Sigma_+\pm(1+\Sigma_+)\tilde{h}|},$$

$$-1 < \Sigma_+ < \min\left(-\frac{1\pm 2\tilde{h}}{4\pm 2\tilde{h}}, -\frac{\tilde{h}^2}{3+\tilde{h}^2}\right).$$

Here the situation is more complicated than before. However, from the eigenvalues we see that the equilibrium points $\tilde{\mathcal{F}}_+(h)$ and $\tilde{\mathcal{E}}_+(h)$ always have an unstable mode. Considering the eigenvalues for the remaining equilibrium points we note that there are no stable equilibrium points for $h > -1$ ($\tilde{h} > 1$). For $h < -1$ ($\tilde{h} < 1$), there are always some stable plane-wave equilibrium points except in a tiny region. This tiny region is the corresponding type VI_h loophole. As in the case of type IV, in the loophole there seems to be a closed curve, $\mathcal{C}(h)$, acting as an attractor. The curve $\mathcal{C}(h)$ lies in the invariant subspace $y = \chi\sqrt{1-\lambda}/\sqrt{2}$, $z = -\chi\sqrt{1+\lambda}/\sqrt{2}$, along with all of the other attractors. An illustration from the loophole is given in Figure 3.




FIGURE 3. To the left a magnification of the type VI_h loop-hole for the plane waves (where $\Sigma_+ = -\tilde{h}^2/(3 + \tilde{h}^2)$) is presented. The vertical line is $\tilde{h} = \sqrt{2} - 1$, which marks the transition of stability between $\tilde{\mathcal{L}}_-(h)$ and $\tilde{\mathcal{E}}_-(h)$. The right picture shows the mussel attractor for $\tilde{h} = 0.41$, $\gamma = 1.225$. Two solution curves are plotted; one approaches $\mathcal{C}$ from the inside, while the other approaches $\mathcal{C}$ from the outside.




FIGURE 4. A plot of the regions of stability of the various equilibrium points. In the above plot we have set Σ_+ equal to the threshold value $\frac{1}{3h-1}$. Note that there are no stable plane-waves for $h > -1$. The loop-hole is approximately where the line $h = -3 - 2\sqrt{2}$ ($\tilde{h} = \sqrt{2} - 1$) terminates. The question marks indicate regions where no stable equilibrium points are presently known.

By a careful analysis of the remaining equilibrium points and their eigenvalues, we obtain the following result:

- For $h < -1$ and $\gamma > \frac{2(1-h)}{1-3h}$, there are always future stable plane-wave solutions.

- For $h > -1$ and $0 < \gamma < 2$, there are no future stable vacuum plane-wave equilibrium points.

We note that the critical unstable mode at $h = -1$ ($\tilde{h} = 1$) is the matter density Ω . Furthermore, we know that there are no other stable equilibrium points with $\Omega = 0$ (for $h \neq -1/9$). This is consequently a signal that for $h > -1$, the future asymptote is non-vacuum.

In [15] the non-tilted Collins perfect fluid solutions of type VI_h were investigated. This analysis showed that for $h > -1$, the Collins solutions were unstable against tilt whenever $\gamma > \frac{2(3+\sqrt{-h})}{5+3\sqrt{-h}}$. This threshold value is the line separating the regions marked “Collins perfect fluid” and those with a question mark in Figure 4. This instability indicates that for $\gamma > \frac{2(3+\sqrt{-h})}{5+3\sqrt{-h}}$ the late-time asymptote is tilted; however, no stable tilted equilibrium points are presently known in this region.

5.4. The exceptional case: The Collinson-French vacuum. The plane-wave-type analysis above can also be applied to the exceptional case of Bianchi type $VI_{-1/9}$. The non-tilted analysis showed that the Collinson-French vacuum (or the Robinson-Trautman solution) is stable in the non-tilted Bianchi cosmologies for $\gamma > 10/9$ [26]. However, the tilted analysis showed that there is one unstable mode for $\gamma > 4/3$ and one more for $\gamma > 14/9$ [6, 15]. Hence, a tilted analysis is required to check the stability.

The Collinson-French solution is given by (in the N-gauge):

$$(5.8) \quad \begin{aligned} \Sigma_+ &= -\frac{1}{3}, \quad \Sigma_- = \frac{1}{3\sqrt{3}}, \quad \Sigma_{13} = \frac{\sqrt{5}}{3\sqrt{3}}, \quad N_{23} = \frac{1}{\sqrt{2}}, \quad A = \frac{1}{\sqrt{6}}, \\ \Sigma_{12} &= \Sigma_{23} = \Omega = \lambda = 0. \end{aligned}$$

In this case the v_i -equations also decouple and we can again treat these separately. For the exceptional case, $h = -1/9$, which leads to the exact vanishing of one of the constraint equations. Hence, the tilt-velocity can only have 2 independent components and we set $v_3 = 0$.

Writing $(X, Y) = (v_1, v_2)$, we obtain

$$(5.9) \quad \begin{aligned} X' &= \left(T - \frac{2}{3}\right)X - \frac{2}{3}\sqrt{6}Y^2, \\ Y' &= \left(T + \frac{2}{3}\sqrt{6}X\right)Y, \end{aligned}$$

where T is defined as usual with

$$V^2 S = \frac{2}{3}X^2.$$

The above system possesses the following equilibrium points:

- (1) $\mathcal{CF}$: $X = Y = 0$, $0 < \gamma < 2$.
- (2) $\widetilde{\mathcal{CF}}_{1\pm}$: $X = -\frac{\sqrt{6}(3\gamma-4)}{2(3-\gamma)}$, $Y = \pm \frac{\sqrt{5(3\gamma-4)(3-2\gamma)}}{\sqrt{2}(3-\gamma)}$, $\frac{4}{3} < \gamma < \frac{3}{2}$.
- (3) $\widetilde{\mathcal{CF}}_2$: $X = \frac{\sqrt{6}(9\gamma-14)}{6(\gamma-1)}$, $Y = 0$, $\frac{24-\sqrt{6}}{15} < \gamma < \frac{24+\sqrt{6}}{15}$.
- (4) $\widetilde{\mathcal{ECF}}_{\pm}$: $X = \pm 1$, $Y = 0$, $0 < \gamma < 2$.

We note that $Y = 0$ is an invariant subspace of the system, which splits the unit disc, D^2 , into two. These two halves are related via the symmetry transformation $Y \mapsto -Y$. Hence, it suffices to analyse the system (5.9) for $Y \geq 0$.




FIGURE 5. Solution curves and phase portraits for the dynamical system (5.9). From top left to right: $\gamma = 1.2$, $\gamma = 1.4$, $\gamma = 1.46$, $\gamma = 1.48$, $\gamma = 1.6$, and $\gamma = 1.8$.

Based on the eigenvalues of the equilibrium points, the equilibrium points are future attractors in the range specified:

- (1) $\mathcal{CF}$: $10/9 < \gamma \leq 4/3$.
- (2) $\widetilde{\mathcal{CF}}_{1\pm}$: $\frac{4}{3} < \gamma < \frac{5}{6} + \frac{\sqrt{721}}{42}$.
- (3) $\widetilde{\mathcal{CF}}_2$: Never an attractor.
- (4) $\widetilde{\mathcal{ECF}}_-$: $\frac{24-\sqrt{3}}{15} < \gamma < 2$.

In Figure 5 we have plotted some phase portraits for different values of γ .

Including the previous analysis of the non-tilted equilibrium points we can thus conclude:

- $2/3 < \gamma < 10/9$: Collins type $\text{VI}_{-1/9}$ perfect fluid solution is stable.
- $\gamma = 10/9$: Wainwright's type $\text{VI}_{-1/9}^*$, $\gamma = 10/9$, solution [27] is stable.
- $10/9 < \gamma < 2$: The Collinson-French vacuum is stable. Moreover, for $4/3 < \gamma$ the asymptotic tilt velocity is non-zero and for $\frac{5}{6} + \frac{\sqrt{721}}{42} < \gamma$ it is extreme.

6. SUMMARY AND OUTLOOK

In this paper we have considered tilted Bianchi models of solvable type. We have used the dynamical systems approach and written down the system of equations with one unspecified gauge function. We pointed out that various different gauge choices are appropriate for different models and different applications. This formalism was then used to study the late-time behaviour of various tilted Bianchi

Bianchi Type	Matter	Attractor	Comments
I	$\frac{2}{3} < \gamma < 2$	$\mathcal{I}(I)$	No tilt allowed
II	$\frac{2}{3} < \gamma < \frac{10}{7}$ $\frac{10}{7} < \gamma < \frac{14}{9}$ $\gamma = \frac{14}{9}$ $\frac{14}{9} < \gamma < 2$	$\mathcal{CS}(II)$ $\mathcal{H}(II)$ $\mathcal{L}(II)$ $\mathcal{E}(II)$	Non-tilted Collins-Stewart Hewitt's tilted type II [28] Tilted type II bifurcation Extremely tilted
IV	$\frac{2}{3} < \gamma < 2$	Plane waves	Tilted/non-tilted
V	$\frac{2}{3} < \gamma < 2$	Milne	Tilted/non-tilted
$VI_{h < -1}$	$\frac{2}{3} < \gamma < \frac{2(1-h)}{1-3h}$ $\frac{2(1-h)}{1-3h} < \gamma < 2$	$\mathcal{C}(VI_h)$ Plane waves	Non-tilted Collins type VI_h Tilted/non-tilted
$VI_{h > -1}$	$\frac{2}{3} < \gamma < \frac{2(3+\sqrt{-h})}{5+3\sqrt{-h}}$ $\frac{2(3+\sqrt{-h})}{5+3\sqrt{-h}} < \gamma < 2$	$\mathcal{C}(VI_h)$?	Non-tilted Collins type VI_h No attractors known
$VI_{-1/9}^*$	$\frac{2}{3} < \gamma < \frac{10}{9}$ $\gamma = \frac{10}{9}$ $\frac{10}{9} < \gamma < 2$	$\mathcal{C}(VI_{-1/9})$ $\mathcal{W}(VI_{-1/9})$ Collinson-French	Non-tilted Collins type $VI_{-1/9}$ Non-tilted Wainwright type $VI_{-1/9}$ Tilted/non-tilted
VI_0	$\frac{2}{3} < \gamma < \frac{6}{5}$ $\gamma = \frac{6}{5}$ $\frac{6}{5} < \gamma < 2$	$\mathcal{C}(VI_0)$ $\mathcal{L}(VI_0)$ $\mathcal{E}(VI_0)$	Non-tilted Collins type VI_0 Tilted type VI_0 bifurcation Extremely tilted
VII_h	$\frac{2}{3} < \gamma < 2$	Plane waves	Tilted/non-tilted
VII_0	$\frac{2}{3} < \gamma < 2$	?	Analysis incomplete

TABLE 1. The late-time behaviour of Bianchi models with a tilted γ -law perfect fluid (see the text for details and references). The case $0 < \gamma < 2/3$ is covered by Corollary 3.6.

models, with an emphasis on the existence of equilibrium points and their stability properties.

Of particular interest are the plane-wave spacetimes of the class B models. In the analysis we have seen a new feature arise in the dynamical behaviour of Bianchi cosmologies; namely, an oscillating behaviour of the asymptotic tilt velocity. In the type IV model, this oscillation occurs in a tiny region of the parameter space in which the dynamical system undergoes a so-called Hopf-bifurcation, and is related to the fact that there are no stable equilibrium points for the corresponding range of parameters in the compact state space. This Hopf-bifurcation is a manifestation of a more general feature of the asymptotic behaviour of these models; their extreme sensitivity to the parameter values. An outline of the stability properties of the plane-waves was given and we pointed out that they are future attractors in the set of type IV and VII_h tilted Bianchi models. These models are studied further, both numerically and analytically, in a companion paper [18].

The plane-wave analysis in the type IV models has emphasised that studying the equilibrium points alone is not sufficient to determine their late-time behaviour. This becomes even more manifest for the type VII_h models where more attracting closed orbits appear. In addition, numerical analysis suggests that for the type VII_h analogue of the loophole, the solution curves will approach a compact surface which is topologically a torus [18]. As noted earlier, these interesting dynamical

results are not gauge effects; however, since the interesting mathematical effects occur in the $\Omega = 0$ invariant set, it is not clear what the physical implications of this behaviour is.

In a companion paper the tilted type IV and VII_h models are studied in greater detail, both analytically and numerically [18]. As in this paper, we will be mostly interested in local behaviour close to equilibrium points, although we will also consider some global aspects. In particular, a more detailed analysis of the closed curves is performed. This analysis confirms the claims in this paper.

The late-time behaviour of tilted Bianchi models is summarized in Table 1. This Table is based on the results in this paper, and the following references: Type II [14]; Type V [13]; Type VI_h [15]; Type $VI_{-1/9}^*$ [15]; Type VI_0 [16]. As can be seen from this table, there is a gap in our knowledge and it is of interest to analyze the late-time behaviour of Bianchi type VI_h and type VII_0 models. Both of these models seem to have a different type of behaviour to that of the other Bianchi models studied here; the type VI_h model possibly has a matter-dominated late-time behaviour and the type VII_0 model might have, in the terminology of [29], an *extreme Weyl dominance* at late times. Here we have seen that this latter behaviour is also present in the one-component tilted type VII_0 subspace $T_1(VII_0)$. The remaining Bianchi models (which are excluded in this work) are the two semi-simple models of type VIII and type IX. Both of these models need a different approach than the one adopted here.

We have focused on the late-time behaviour in this work. An equally interesting task would be to consider the early time behaviour. In [14] it was argued that the tilted type II is *chaotic* initially. We note that all Bianchi types, except for type I and V, have type II as part of the boundary. For the tilted type VI_0 model [16] the type II boundary played a crucial role in the initial singular behaviour, and since the type II boundary is chaotic, it was argued that the tilted type VI_0 is chaotic as well. We might therefore wonder whether a similar phenomenon occurs in all the tilted Bianchi models having type II as part of the boundary. Indeed, since type V does not have the type II as part its boundary, we can ask whether tilted Bianchi type V model is the only tilted Bianchi model (apart from the trivial type I) which does not have chaotic behaviour initially. A combined dynamical systems and metric approach [30] and a Hamiltonian analysis [31] suggest that this is indeed the case. Moreover, there is some numerical evidence for this in the current tilted Bianchi models under consideration. We shall return to this question in future work.

7. ACKNOWLEDGMENTS

The authors would like to thank Robert van den Hoogen for discussions. This work was supported by NSERC (AC) and the Killam Trust and AARMS (SH).

APPENDIX A. CONSISTENCY OF THE CONSTRAINTS

We define the following functions:

$$\begin{aligned}
 H &= \Sigma^2 + A^2 + |\mathbf{N}_\times|^2 + \Omega - 1 \\
 Q_1 &= 2\Sigma_+ A + 2\text{Im}(\Sigma_\times^* \mathbf{N}_\times) + \frac{\gamma\Omega v_1}{G_+} \\
 \mathbf{Q} &= \Sigma_1(i\bar{N} - \sqrt{3}A) + i\Sigma_1^* \mathbf{N}_\times + \frac{\gamma\Omega \mathbf{v}}{G_+} \\
 G &= A^2 + 3h(|\mathbf{N}_\times|^2 - \bar{N}^2).
 \end{aligned}
 \tag{A.1}$$

The constraints are satisfied if and only if all of these functions vanish. To check the consistency of these constraints, we assume first that they do not vanish and calculate their time-derivatives. After some algebra we obtain

$$\begin{aligned}
 H' &= 2qH + 2AQ_1, \\
 Q_1' &= 2(q - 1 + \Sigma_+)Q_1 - 2\sqrt{3}\text{Re}(\Sigma_1 \mathbf{Q}^*), \\
 \mathbf{Q}' &= (2q - 2 - \Sigma_+ + i\phi')\mathbf{Q} - \sqrt{3}\Sigma_\times \mathbf{Q}^*, \\
 G' &= 2(q + 2\Sigma_+)G.
 \end{aligned}
 \tag{A.2}$$

This means that if all the constraints vanish initially, they will be zero at all times.

APPENDIX B. SOME PROOFS

B.1. Outline of the proof of Theorem 3.3. We will prove Theorem 3.3 for the Bianchi type VII_h models by contradiction. We assume, therefore, that there exists an equilibrium point with $\mathbf{v} \neq 0$ and $\bar{N}^2 - |\mathbf{N}_\times|^2 > 0$ and $A \neq 0$. The outline of the proof goes as follows.

We first assume that $\mathbf{N}_\times = 0$. This leads to

$$\bar{N} \neq 0 \Rightarrow \Sigma_\times = 0 \Rightarrow \Sigma_1^2 = -\frac{\gamma\Omega}{2G_+} \mathbf{v}^2.$$

Using the constraint equation we see that $\Omega \neq 0$ and $(i\bar{N} - \sqrt{3}A)^2$ equals a negative real number. Thus $A = 0$, which is a contradiction. We can thus assume that $\mathbf{N}_\times \neq 0$. This allows us to use the N-gauge, and we will assume this gauge is chosen henceforth.

Let us now assume that $\Omega \neq 0$. Using the evolution equations for A , $\mathbf{N}_\times$, Σ_1 , and $\mathbf{v}$ we obtain $\Sigma_{23} = 0$, $T = \mathcal{S}$, $q = 2 + 3\mathcal{S}$, $Av_1 = -1 - (5/2)\mathcal{S}$, $\Sigma_+ = -1 - (3/2)\mathcal{S}$, and $\Sigma_- = v_1 N_{23}$. Inserting this information into the Ω equation we obtain the two separate cases: (i) $V^2 = 1$, and (ii) $\gamma = 6/5$. After some lengthy calculations both of these cases lead to contradictions. These calculations involve solving the remaining equations and making sure that all parameters are real and obey the required bounds; i.e., $0 < \Omega < 1$, $0 < V^2 \leq 1$, etc.

Therefore, we must have $\Omega = 0$. In this case the equations of motion decouple and we can use the non-tilted analysis which implies that the only equilibrium points are the plane-wave spacetime. From eq.(5.7) we require $\varphi' = 0$, which implies $x = 1$ ($\lambda^2 > 1$). Since $x = v_1$, this implies that $V^2 = v_1^2 + v_2^2 + v_3^2 \geq 1$. However, since $V^2 \leq 1$, we must have $V^2 = 1$ and $v_2 = v_3 = 0$ which is a contradiction. Since $\mathbf{v} = 0$ implies $\Sigma_1 = 0$, Theorem 3.3 now follows.

B.2. Proof of Theorem 4.3. The Proof consists of two steps: (i): The existence of a monotonic function. (ii): Showing that the monotonic function forces $\Sigma_+ \rightarrow -(3\gamma - 4)/2$ at late times.

The first step is to find a monotonic function for the reduced system and amend it so that it is monotonic for the full system at late times. Such a function is:

$$(B.1) \quad \hat{Z}_4 = \frac{\sigma_1^m \beta \Omega}{(1 + m\Sigma_+)^{2(1+m)}}, \quad m \equiv \frac{1}{2}(3\gamma - 4),$$

where β is given in eq.(4.3). We amend this function by defining

$$\bar{Z}_4 = \frac{\hat{Z}_4}{1 + fM \cos \psi}, \quad f \equiv \frac{\rho [\sigma_1(1 - m^2) - m(\Sigma_+ + 1)(1 + m\Sigma_+)]}{2\sqrt{3}\sigma_1(1 + m\Sigma_+)}.$$

Note that since $|\rho/\sigma_1| \leq 1$ and $m \leq 1$, f is bounded. At sufficiently late times we have

$$(B.2) \quad \frac{\bar{Z}_4'}{\bar{Z}_4} = \frac{2(\Sigma_+ + m)^2}{(1 + m\Sigma_+)} + MB,$$

where B is a function of the state space variables and is bounded. Thus at late times, $\bar{Z}_4$ is monotonically increasing. However, for $4/3 < \gamma < 2$, this function is bounded for sufficiently late times, and hence, we get the bound

$$(B.3) \quad \int_{\tau_0}^{\infty} (\Sigma_+ + m)^2 d\tau < \infty,$$

for a sufficiently large τ_0 .

The second step is to show that this implies $\Sigma_+ \rightarrow -m$. We will show this by contradiction.

Define the everywhere positive function $f(\tau) \equiv (\Sigma_+ + m)^2$. Assume that $f \not\rightarrow 0$. Then there must exist a $\delta > 0$ and an infinite sequence $\{\tau_n\}$ with $\lim_{n \rightarrow \infty} \tau_n = \infty$ such that

$$(B.4) \quad f(\tau_n) \geq \delta, \quad \forall n.$$

From the evolution equations we note that $f'(\tau)$ is bounded; i.e. there exists a $D > 0$ such that $|f'(\tau)| \leq D$ for all τ . We then define the family of functions g_n given by

$$(B.5) \quad g_n(\tau) \equiv \begin{cases} \delta - D|\tau_n - \tau|, & \tau_n - \delta/D \leq \tau \leq \tau_n + \delta/D \\ 0, & \text{otherwise.} \end{cases}$$

We note that $0 \leq g_n(\tau) \leq f(\tau)$, and $\int_{\tau_0}^{\infty} g_n(\tau) d\tau = \delta^2/D$. Moreover, by defining (we can without loss of generality assume that $\text{support}(g_i) \cap \text{support}(g_j) = \emptyset$ for $i \neq j$)

$$g(\tau) \equiv \sum_n^{\infty} g_n(\tau),$$

we have $0 \leq g(\tau) \leq f(\tau)$. Thus

$$(B.6) \quad \int_{\tau_0}^{\infty} f(\tau) d\tau \geq \int_{\tau_0}^{\infty} g(\tau) d\tau = \sum_n^{\infty} \int_{\tau_0}^{\infty} g_n(\tau) d\tau = \frac{\delta^2}{D} \sum_n^{\infty} 1 = \infty,$$

which contradicts eq.(B.3). Hence, $\Sigma_+ \rightarrow -m$ which proves the Theorem.

APPENDIX C. EIGENVALUES FOR NON-EXCEPTIONAL TYPES.

Below, we have defined $\tilde{h} = 1/\sqrt{|h|}$ and

$$(C.1) \quad \beta = \begin{cases} i\tilde{h}, & h > 0, \\ 0, & \text{IV}, \\ \tilde{h}, & h < 0. \end{cases}$$

C.1. Plane waves, $\mathcal{L}(h)$ (for all h):

$$\begin{aligned} \lambda_1 &= 0, \quad \lambda_{2,3} = -2 \left[(1 + \Sigma_+) \pm 2i\sqrt{3}N_{23} \right], \quad \lambda_4 = -4\Sigma_+ - (3\gamma - 2), \\ \lambda_5 &= (3\gamma - 4) + 2\Sigma_+, \quad \lambda_{6,7} = (3\gamma - 4) - \Sigma_+ \pm (1 + \Sigma_+)\beta. \end{aligned}$$

C.2. Plane waves, $\tilde{\mathcal{L}}(h)$ (for all h):

$$\begin{aligned} \lambda_1 &= 0, \quad \lambda_{2,3} = -2 \left[(1 + \Sigma_+) \pm 2i\sqrt{3}N_{23} \right], \quad \lambda_4 = -\frac{2-\gamma}{\gamma-1}(1 - 2\Sigma_+), \\ \lambda_5 &= -\frac{(1 - 2\Sigma_+)(5\gamma - 6 + 2\gamma\Sigma_+)(3\gamma - 4 + 2\Sigma_+)}{(\gamma - 1)(9\gamma - 10 + 4\Sigma_+(1 - \Sigma_+))}, \\ \lambda_{6,7} &= \frac{(1 - 2\Sigma_+)(3\gamma - 4)}{2(\gamma - 1)} \pm (1 - v_1)(1 + \Sigma_+)\beta. \end{aligned}$$

C.3. Plane waves, $\tilde{\mathcal{L}}_+(h)$ (for all h):

$$(C.2) \quad \begin{aligned} \lambda_1 &= 0, \quad \lambda_{2,3} = -2 \left[(1 + \Sigma_+) \pm 2i\sqrt{3}N_{23} \right], \quad \lambda_4 = 0, \\ \lambda_5 &= 2(1 - 2\Sigma_+), \quad \lambda_{6,7} = 1 - 2\Sigma_+. \end{aligned}$$

C.4. Plane waves, $\tilde{\mathcal{L}}_-(h)$ (for all h):

$$\begin{aligned} \lambda_1 &= 0, \quad \lambda_{2,3} = -2 \left[(1 + \Sigma_+) \pm 2i\sqrt{3}N_{23} \right], \quad \lambda_4 = -4(1 + \Sigma_+), \\ \lambda_5 &= -\frac{2(5\gamma - 6 + 2\gamma\Sigma_+)}{2 - \gamma}, \quad \lambda_{6,7} = -1 - 4\Sigma_+ \pm 2(1 + \Sigma_+)\beta. \end{aligned}$$

C.5. Plane waves, $\tilde{\mathcal{F}}_{\pm}(\text{IV})$:

$$\begin{aligned} \lambda_1 &= 0, \quad \lambda_{2,3} = -2 \left[(1 + \Sigma_+) \pm 2i\sqrt{3}N_{23} \right], \quad \lambda_4 = -\frac{(1 - 2\Sigma_+)(5\gamma - 6)}{3 - 2\gamma}, \\ \lambda_5 &= 0, \quad \lambda_6\lambda_7 = -\frac{(1 - 2\Sigma_+)^2(4 - 3\gamma)(3\gamma - 4 - \Sigma_+)(3 - 2\gamma + \gamma\Sigma_+)}{(3 - 2\gamma)G(\gamma, \Sigma_+)} \\ \lambda_6 + \lambda_7 &= \frac{(1 - 2\Sigma_+)F(\gamma, \Sigma_+)}{4(3 - 2\gamma)(17\gamma^2 - 40\gamma + 24)G(\gamma, \Sigma_+)} \end{aligned}$$

where

$$(C.3) \quad \begin{aligned} G(\gamma, \Sigma_+) &\equiv (5\gamma - 6)\Sigma_+^2 - (18 - 25\gamma + 9\gamma^2)\Sigma_+ - 3 + 2\gamma, \\ F(\gamma, \Sigma_+) &\equiv \left[2(17\gamma^2 - 40\gamma + 24)\Sigma_+ - 33\gamma^3 + 121\gamma^2 - 152\gamma + 66 \right]^2 \\ &\quad - 9(\gamma - 1)^2 (121\gamma^4 - 736\gamma^3 + 1664\gamma^2 - 1656\gamma + 612) \end{aligned}$$

Here is $G(\gamma, \Sigma_+) < 0$ in the whole region under consideration. $F(\gamma, \Sigma_+) = 0$, defines a line from $(4/3, 0)$ to $(4/3, -1/4)$.

C.6. Plane waves, $\tilde{\mathcal{E}}_{\pm}(IV)$:

$$\begin{aligned}\lambda_1 &= 0, & \lambda_{2,3} &= -2 \left[(1 + \Sigma_+) \pm 2i\sqrt{3}N_{23} \right], & \lambda_4 &= \frac{(1 - 2\Sigma_+)(1 + 2\Sigma_+)}{\Sigma_+}, \\ \lambda_5 &= 0, & \lambda_6 &= -\frac{(1 - 2\Sigma_+)(1 + 4\Sigma_+)}{3\Sigma_+}, & \lambda_7 &= \frac{2(1 - 2\Sigma_+)(2\gamma - 3 - \gamma\Sigma_+)}{3\Sigma_+(2 - \gamma)},\end{aligned}$$

C.7. Plane waves, $\tilde{\mathcal{F}}_{\pm}(h)$:

$$\begin{aligned}\lambda_1 &= 0, & \lambda_{2,3} &= -2 \left[(1 + \Sigma_+) \pm 2i\sqrt{3}N_{23} \right], \\ \lambda_4 &= -\frac{(1 - 2\Sigma_+)[(5\gamma - 6) \pm \tilde{h}(2 - \gamma)]}{3 - 2\gamma \pm \tilde{h}}, & \lambda_5 &= 2(1 - v_1)\alpha, \\ \lambda_6\lambda_7 &\propto (1 - 2\Sigma_+)^2[(4 - 3\gamma)(1 + \Sigma_+) + \alpha(2 - \gamma)] \\ &\quad \times (3\gamma - 4 - \Sigma_+ - \alpha)[3 - 2\gamma + \gamma(\Sigma_+ + \alpha)] \\ \text{(C.4)} \quad \lambda_6 + \lambda_7 &= \text{something really awful}\end{aligned}$$

where $\alpha \equiv \pm(1 + \Sigma_+)\tilde{h}$. The expressions for λ_6 and λ_7 are too nasty to write down, but $\lambda_6 + \lambda_7 = 0$ defines a curve which ultimately gives rise to the type VI_h loopole.

C.8. Plane waves, $\tilde{\mathcal{E}}_{\pm}(h)$:

$$\begin{aligned}\lambda_1 &= 0, & \lambda_{2,3} &= -2 \left[(1 + \Sigma_+) \pm 2i\sqrt{3}N_{23} \right], & \lambda_4 &= \frac{(1 - 2\Sigma_+)(3 + 6\Sigma_+ + 2\alpha)}{3\Sigma_+ + \alpha}, \\ \lambda_5 &= 2(1 - v_1)\alpha, & \lambda_6 &= -\frac{(1 - 2\Sigma_+)(1 + 4\Sigma_+ + 2\alpha)}{3\Sigma_+ + \alpha}, \\ \lambda_7 &= \frac{2(1 - 2\Sigma_+)[2\gamma - 3 - \gamma(\Sigma_+ + \alpha)]}{(3\Sigma_+ + \alpha)(2 - \gamma)},\end{aligned}$$

where $\alpha \equiv \pm(1 + \Sigma_+)\tilde{h}$.

APPENDIX D. EIGENVALUES FOR THE COLLINSON-FRENCH VACUUM

Below are listed the eigenvalues corresponding to the eq. (5.9), $\lambda_{1,2}$, and the eigenvalue corresponding to the matter density, Ω , which is denoted λ_3 .

D.1. Collinson-French, $\mathcal{CF}$:

$$\lambda_1 = \frac{1}{3}(9\gamma - 14), \quad \lambda_2 = (3\gamma - 4), \quad \lambda_3 = -\frac{1}{3}(9\gamma - 10).$$

D.2. Collinson-French, $\widetilde{\mathcal{CF}}_{1\pm}$:

$$\lambda_{1,2} = \frac{s \pm \sqrt{s^2 - 4d}}{2}, \quad \lambda_3 = -\frac{10\gamma(4\gamma^2 - 9\gamma + 6)}{3(3 - \gamma)(3\gamma^2 - 9\gamma + 10)}$$

where

$$d \equiv \lambda_1\lambda_2 = \frac{100(3\gamma - 4)(3 - 2\gamma)}{3(3 - \gamma)(3\gamma + 1)}, \quad s \equiv \lambda_1 + \lambda_2 = \frac{105 \left[\left(\gamma - \frac{5}{6} \right)^2 - \frac{721}{42^2} \right]}{3(3 - \gamma)(3\gamma + 1)}.$$

D.3. Collinson-French, $\widetilde{\mathcal{CF}}_2$:

$$\begin{aligned}\lambda_1 &= -\frac{10(3-2\gamma)}{3(\gamma-1)}, & \lambda_2 &= \frac{(9\gamma-14)[(9\gamma-14)^2-6(\gamma-1)^2]}{3(81\gamma-258\gamma+202)}, \\ \lambda_3 &= -\frac{10\gamma(12\gamma^2-39\gamma+32)}{3(\gamma-1)(81\gamma^2-246\gamma+190)}.\end{aligned}\tag{D.1}$$

D.4. Collinson-French, $\widetilde{\mathcal{ECF}}_{\pm}$:

$$\lambda_1 = \frac{2}{3}(1 \pm \sqrt{6}), \quad \lambda_2 = -\frac{2[9\gamma-14 \mp \sqrt{6}(\gamma-1)]}{3(2-\gamma)}, \quad \lambda_3 = -\frac{1}{3}(4 \mp \sqrt{6}).$$

REFERENCES

1. A.R. King and G.F.R. Ellis, *Commun. Math. Phys.* **31** (1973) 209
2. G.F.R. Ellis and M.A.H. MacCallum, *Comm. Math. Phys.* **12** (1969) 108
3. C.G. Hewitt and J. Wainwright in *Dynamical Systems in Cosmology*, eds: J. Wainwright and G.F.R. Ellis, Cambridge University Press (1997)
4. A.A. Coley, *Dynamical Systems and Cosmology*, Kluwer, Academic Publishers (2003)
5. V.A. Belinskii, I.M. Khalatnikov and E.M. Lifshitz, *Adv. Phys.* **31** (1982) 639
6. J.D. Barrow and D.H. Sonoda, *Phys. Reports* **139** (1986) 1
7. K. Rosquist and R. T. Jantzen, *Phys. Rep.* **166** (1988) 89
8. O.I. Bogoyavlenskii and S.P. Novikov, *Sov. Math. Sov.* **2** (1982) 159; originally published as *Trudy Sem. Petrovsk.* **1** (1975) 7; Bogoyavlenskii, O. I. (1985) *Methods in the Qualitative Theory of Dynamical Systems in Astrophysics and Gas Dynamics* Springer-Verlag.
9. J.D. Barrow and D.H. Sonoda, *Gen. Rel. Grav.* **17** (1985) 409
10. I.S. Shikin, *Sov. Phys. JETP* **41** (1976) 794
11. C.B. Collins, *Comm. Math. Phys.* **39** (1974) 131
12. C. B. Collins and G. F. R. Ellis, *Phys. Rep.* **56** (1979) 65
13. C.G. Hewitt and J. Wainwright, *Phys. Rev.* **D46** (1992) 4242
14. C.G. Hewitt, R. Bridson, J. Wainwright, *Gen. Rel. Grav.* **33** (2001) 65
15. J.D. Barrow and S. Hervik, *Class. Quantum Grav.* **20** (2003) 2841
16. S. Hervik, *Class. Quantum Grav.* **21** (2004) 2301
17. A. Coley and S. Hervik, *Class. Quantum Grav.* **21** (2004) 4193-4208
18. S. Hervik, R. van den Hoogen and A. Coley, *Future asymptotic behaviour of tilted Bianchi models of type IV and VII_h*.
19. H. van Elst, *Hubble normalised orthonormal frame equations for perfect fluid cosmologies in component form*, available at www.maths.qmul.ac.uk/~hve/research.html.
20. C.G. Hewitt and J. Wainwright, *Class. Quant. Grav.* **10** (1993) 99
21. P. Apostolopoulos, [gr-qc/0407040](https://arxiv.org/abs/gr-qc/0407040)
22. R.M. Wald, *Phys. Rev.* **D28** (1983) 2118
23. J. Wainwright, M.J. Hancock and C. Uggla, *Class. Quant. Grav.* **16** (1999) 2577
24. U.S. Nilsson, M.J. Hancock and J. Wainwright, *Class. Quant. Grav.* **17** (2000) 3119
25. W.C. Lim, *private communication*
26. C.G. Hewitt, J.T. Horwood, J. Wainwright, *Class. Quant. Grav.* **20** (2003) 1743
27. J. Wainwright, *Gen. Rel. Grav.* **16** (1984) 657
28. C.G. Hewitt, *Gen. Rel. Grav* **23** (1991) 691
29. J.D. Barrow and S. Hervik, *Class. Quantum Grav.* **19** (2002) 5173
30. A.A. Peresetsky, in *Topics in Modern Mathematics*, Petrovskii Seminar no 5, ed. O.A. Oleinik, Consultants Bureau (1985)
31. R.T. Jantzen, pages 61-147 in *Proc. Int. Sch. Phys "E. Fermi" Course LXXXVI*, eds. Ruffini and Melchiorri, North Holland, 1987 [[gr-qc/0102035](https://arxiv.org/abs/gr-qc/0102035)]

DEPARTMENT OF MATHEMATICS & STATISTICS, DALHOUSIE UNIVERSITY, HALIFAX, NOVA SCOTIA, CANADA B3H 3J5

E-mail address: `aac@mathstat.dal.ca`, `herviks@mathstat.dal.ca`